

Cytoprotective effect of alpha-2-macroglobulin against pesticide-induced generation of ROS in neuronal SH-SY5Y cells

Swati Dixit¹, Haseeb Ahsan^{2,*}, Fahim Halim Khan^{1,*}



Use your smartphone to scan this QR code and download this article

ABSTRACT

Introduction: Many clinical studies have demonstrated that continuous exposure to pesticides, especially organophosphates and pyrethroids, causes toxicities such as carcinogenicity and neurotoxicity that lead to disorders such as diabetes, lung cancer, and neurodegenerative diseases. The mechanism underlying pesticide-induced neurotoxicity involves the production of ROS, which causes neuronal injury through oxidative stress. **Methods:** In the present study, the neuronal SH-SY5Y cell line was used to investigate the effect of the pesticides chlorpyrifos (organophosphate), aldicarb (carbamate), and deltamethrin (pyrethroid) on ROS-mediated toxicity and the protective effect of alpha-2-macroglobulin (α_2M), a protease inhibitor and beta-amyloid plaque scavenger in the human brain. For cell viability and cytotoxicity, the MTT assay was performed. To monitor ROS production, assays such as DCFHDA, H₂O₂, and MDA were performed, along with assays of the activity of antioxidant enzymes such as superoxide dismutase and glutathione peroxidase. **Results:** The experimental findings suggest a cytoprotective role for α_2M in ROS-mediated toxicity that causes neuronal injury in humans. **Conclusion:** Hence, α_2M could be possibly used as a protective agent against oxidative neurotoxicity caused by pesticides.

Key words: Alzheimer's disease, SH-SY5Y, reactive oxygen species, neurotoxicity, deltamethrin, chlorpyrifos, aldicarb, alpha-2-macroglobulin

INTRODUCTION

The pesticides chlorpyrifos (CPF), aldicarb (ALD), and deltamethrin (DLM; **Figure 1**) are known to cause both acute and chronic neurotoxicity in humans and animals through the inhibition of cholinesterase (ChE) activity in the synaptic junction of neurons, binding to gamma-aminobutyric acid (GABA)-gated channels by preventing the closure of sodium channels¹. Humans are exposed to these pesticides in a number of ways, such as the consumption of fruits and vegetables contaminated with pesticide sprays and the inhalation of fumigants²⁻⁵. Apart from causing substantial toxicity, these pesticides interfere with normal neuronal processes *in vivo*, leading to oxidative damage and cellular inflammation in the human brain⁶⁻⁸. Some of the modifications caused by these pesticides include oxidative stress⁹, disrupted neurotransmission¹⁰ and neuronal differentiation¹¹, inhibition of replication in neuronal cells¹², cognitive changes, decrease in psychomotor functions¹³, and apoptosis in neuronal cells by their metabolites¹⁴⁻¹⁷. Such changes in neurons may lead to various neurodegenerative diseases such as Alzheimer's disease (AD) and/or Parkinson's disease (PD)¹⁸⁻²⁰.

Alpha-2-macroglobulin (α_2M) is produced in the human brain and keeps a check on reactive oxygen

species (ROS)²¹ and β -amyloid plaque formation, thereby preventing neuronal degeneration and AD progression. α_2M is a serum protease inhibitor that may be involved in AD in mediating the clearance and degradation of A β , a major constituent of β -amyloid deposits^{22,23}. α_2M , an acute-phase protein and a major component of the innate immune system, is a cerebrospinal fluid (CSF) marker of neuronal injury in preclinical AD [22]. An increased concentration of α_2M in the blood is associated with neuronal injury²⁴. Previous studies have shown that α_2M interacts with the above mentioned pesticides^{3,21}, indicating the probable role of α_2M as an unexplored biomarker in pesticide-induced ROS-mediated neurotoxicity²⁵. Hence, in the present study, α_2M was used to study the modulating/impairing effect of pesticide generated ROS on the neuronal SH-SY5Y (NCCS, Pune) cell line.²⁶

METHODS

Materials

All pesticides (CPF, ALD, and DLM) and chemicals (MTT dye, DMSO, pyrogallol, DMEM and Ham's F12 medium, fetal bovine serum [FBS], thiobarbituric acid [TBA], 1% penicillin-streptomycin, dichlorodihydrofluorescein diacetate [DCFH-DA], glutathione

¹Department of Biochemistry, Faculty of Life Sciences, Aligarh Muslim University, Aligarh 202002, India

²Department of Biochemistry, Faculty of Dentistry, Jamia Millia Islamia, New Delhi 110025, India

Correspondence

Haseeb Ahsan, Department of Biochemistry, Faculty of Dentistry, Jamia Millia Islamia, New Delhi 110025, India

Email: drhahsan@gmail.com

Correspondence

Fahim Halim Khan, Department of Biochemistry, Faculty of Life Sciences, Aligarh Muslim University, Aligarh 202002, India

Email: fahimhkh@rediffmail.com

History

- Received: Oct 20, 2022
- Accepted: Jan 26, 2023
- Published: Feb 28, 2023

DOI : 10.15419/bmrat.v10i2.792



Cite this article : Dixit S, Ahsan H, Khan F H. **Cytoprotective effect of alpha-2-macroglobulin against pesticide-induced generation of ROS in neuronal SH-SY5Y cells.** *Sci. Tech. Dev. J.*; 10(2):5537-5544.

Copyright

© VNU-HCM Press. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.




Figure 1: Structure of pesticides a) chlorpyrifos b) aldicarb c) deltamethrin.

[GSH], and NADPH) were purchased from Merck-Sigma (St. Louis, MO, USA). All other reagents used were of analytical standard.

METHODS

Purification and characterization of human α_2M

The α_2M protein was isolated and purified from human blood plasma using ammonium sulfate precipitation followed by gel exclusion chromatography as per the method described previously³. A 5% (w/v) native PAGE was performed²¹ and the gel was stained with Coomassie brilliant blue R-250 (0.15% in 10% acetic acid). The gel was de-stained for 12 h in the de-staining solution (10% acetic acid), and the purified α_2M formed a single band on the gel.

Cell culture

The SH-SY5Y cell line was cultured in a medium containing 1:1 DMEM and Ham's F12 medium, 10% FBS, and 1% penicillin-streptomycin²⁶. The cells were treated with a standard solution of pesticides and α_2M , accordingly, to perform the experiments. Cells were used at 3–7 passages. The cells were divided into five groups based on the treatment with pesticides and proteins to obtain results for various stress markers. Group I was the control group comprising only SH-SY5Y cells under standard conditions (37 °C and 5% CO₂). Group II consisted of SH-SY5Y cells incubated with α_2M . The concentration of protein was 2 μM and the incubation time was 3 h (37 °C). Group III comprised pesticide (CPF, ALD, and DLM)-treated SH-SY5Y cells. Cells were treated with 5 μM of each pesticide (CPF, ALD, and DLM) separately for 3 h under standard conditions (37 °C and 5% CO₂). Group IV was the pesticides (CPF, ALD, and DLM) and α_2M group, in which the SH-SY5Y cells were treated with 5 μM of the pesticides for 3 h and then treated with α_2M for 3 h.

Cell viability and proliferation assay

To determine the cytotoxicity and cell viability of the SH-SY5Y cells, the 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide MTT assay was performed in 96-well plates²⁷. The cells were used at 70–80% confluence. Later, the medium was removed and the cells were treated with pesticides (CPF, ALD, and DLM; 5 μM) for 6 h and later incubated with α_2M for 3 h. Afterward, the cells were treated with MTT solution (5 mg/mL stock solution) at a final concentration of 0.5 mg/mL MTT. The cells were incubated for 4 h at 37 °C in 5% CO₂. Finally, the cells were treated with 100 μL DMSO for 10 min to dissolve the formazan crystals. The absorbance was measured using an ELISA plate reader at 570 nm with a reference wavelength of 630 nm and was directly proportional to the number of viable cells. The experiments were performed in triplicate.

Measurement of ROS

ROS production was measured using DCFH-DA²⁸. The cells were treated with α_2M and the pesticides, following which 10 μM DCFH-DA was added to the medium for 1 h at 37 °C for diffusion into the cells. A multi-detection microplate reader was used for fluorescence measurement. The excitation and emission wavelengths for DCFH-DA were 485 and 535 nm, respectively^{28–30}.

Measurement of malonaldehyde (MDA) levels

To detect the generation of ROS by pesticides in the neuronal cell line, the MDA levels were measured. The MDA levels in SH-SY5Y cells were quantified with the TBA reaction. Thiobarbituric acid reactive substances (TBARS) were measured by comparing the absorption to the standard curve of MDA equivalents generated by the acid-catalyzed hydrolysis of tetramethoxypropane³¹. The absorbance was recorded at 532 nm.

Measurement of superoxide dismutase (SOD) activity

The antioxidant enzyme SOD converts superoxide (O₂^{•-}) into H₂O₂ and O₂, which is converted to water by other enzymes. The measurement was carried out as described previously³². To 80 μL of the cell suspension, 2.82 ml of 0.05 mM tris-succinate buffer was added, and the sample was treated for 30 min in a CO₂ incubator. The reaction was initiated by adding 100 μL of 8 mM pyrogallol solution to each well³². The absorbance was read at 420 nm.

Measurement of glutathione peroxidase (GPx) levels

GPx are cytosolic enzymes that catalyze the conversion of H_2O_2 into H_2O and O_2 and the reduction of peroxide radicals ($ROO\cdot$) into alcohol and oxygen. GPx levels were measured by determining the decrease in absorbance at 340 nm upon the oxidation of NADPH to $NADP^+$ ³³.

Statistically analysis

All experiments were repeated thrice and the data shown is the mean \pm SD. P-value < 0.05 is considered as significant difference.

RESULTS

Cell viability and cell cytotoxicity assay

The MTT assay was performed to assess cell viability and cell proliferation activity, in which the quantity of formazan is directly proportional to the number of viable cells. The SH-SY5Y cells treated with all three pesticides (CPF, ALD, and DLM) in group III showed reduced cell viability ($p < 0.001$; **Figure 1**). Almost no effect on viability was seen for cells treated with α_2M (group II), while the effect of the pesticides appeared to be reversed in cells treated with pesticides after incubation with α_2M (group IV).

Measurement of ROS levels

Figure 3 shows the ROS levels induced by the pesticides as monitored by DCFHDA in the control group of SH-SY5Y cells and all three pesticides in group III ($p < 0.001$). The increased fluorescence in group III as compared to the control group and α_2M group indicates the production of ROS. On the contrary, group IV showed a reduction in the fluorescence intensity as compared to group III, indicating the defensive function of α_2M against ROS production.

Measurement of MDA

Figure 4 shows the TBARS activity in the SH-SY5Y cells in various groups. The highest MDA production was recorded in group III, in which the cells were treated with pesticides. However, group II showed no TBARS production. Group IV (pesticides + cells + α_2M), however, showed a reduction in the TBARS level as compared to group III. The MDA levels were normal in the control group and slightly reduced in the α_2M group ($p < 0.001$). However, MDA levels were significantly decreased in the α_2M + pesticides + cells group ($p < 0.001$) compared to the pesticides group. Hence, the protective effects of α_2M were observed with respect to MDA levels due to pesticide-triggered ROS in SH-SY5Y cells.

Measurement of SOD levels and GPx activity

Figure 5 shows the effect of various treatments on SOD levels in the control and other groups. Group III (pesticides group) showed reduced SOD activity as compared to group I (control) and group II (α_2M ; $p < 0.001$), demonstrating the toxicity triggered by the pesticides. On the other hand, group IV (α_2M + pesticides + SH-SY5Y cells) showed a remarkable increase in SOD and GPx activity levels, indicating the protective effect of α_2M against the pesticides. **Table 1** shows the effect of pesticides on the activity of antioxidant enzymes in the cells. The pesticides group showed reduced GPx activity as compared to the control group I and α_2M group II, while group IV (α_2M and pesticides) showed increased GPx activity levels.

DISCUSSION

CPF, ALD, and DLM are potent neurotoxic pesticides that affect various neuronal processes involved in the synaptic transmission and growth of neurons³⁴⁻³⁷. α_2M is an acute-phase protein produced in the brain to keep ROS levels under control^{3,21} and prevent amyloid plaque formation by dissolving them through receptor-mediated endocytosis. It also protects the brain from undergoing neuronal degeneration²⁵. α_2M is also a major component of the innate immune system and acts as a hallmark of neuronal injury²¹. The human brain synthesizes and secretes α_2M upon stimulation with interleukin-6, which indicates that α_2M is an acute-phase protein in the human CNS. In this study, we characterized ROS formed by CPF, ALD, and DLM in the SH-SY5Y cell line. Furthermore, we investigated the ability of α_2M to modulate the neurotoxic effect of these pesticides. The cellular toxicity of these pesticides on SH-SY5Y cells was first measured by the MTT assay, which indicated the toxic effects of the pesticides on the cell line.

DCFH-DA is a fluorimetric dye that is applied to quantify ROS³⁸. This non-fluorescent compound diffuses into the cells, where it is sequentially hydrolyzed to release DCFH as the substrate by intracellular enzymes. ROS oxidize DCFH and convert it into DCF, a fluorescence product that is measured by spectrofluorometry using a microplate reader. A significant increase in the production of DCF was seen in pesticide-treated cells. This increased production of ROS results in oxidative damage to cellular components. However, when α_2M was incubated with pesticide-treated cells, a contrary effect was observed. Similarly, regarding MDA levels, pesticide-treated groups showed a significant increase in absorbance, but in group IV, a




Figure 2: MTT assay of various groups of SH-SY5Y cells and all three pesticides (CPF, ALD, and DLM). Cells were pretreated with pesticides at 5 μM for 3 h and then incubated with α_2M for 3 h. After incubation, cell viability was evaluated by the MTT assay. The results are shown as the mean \pm SEM of three experiments, with each experiment performed in triplicate ($p < 0.001$ versus control). Group II is not statistically significant to Groups I, III and IV.




Figure 3: DCFHDA ROS assay of SH-SY5Y cells. ROS were determined by the DCFH-DA method. DCFH-DA measurements are reported as fluorescence intensities (AU, arbitrary fluorescence units). Results are mean \pm standard error of triplicates. The error bar depicts the standard error among the six different samples of that group. Cells were incubated with 5 μM of each pesticide for 45 min, and the α_2M concentration was 2 μM . Excitation and emission wavelengths for DCFH-DA were 485 and 535 nm, respectively. Group II is not statistically significant to Groups I, III and IV.




Table 1: Effect of pesticides on the activity of antioxidant enzymes of the groups

Enzyme	Control	CPF	ALD	DLM
SOD	1.6 \pm 0.140	0.90 \pm 0.086	1.4 \pm 0.131	0.88 \pm 0.041
GPx	24.4 \pm 2.25	18.3 \pm 1.31	20 \pm 0.100	16.3 \pm 0.07

SOD and GPx measurements are reported in $\mu\text{mol}/\text{min}/\text{mg}$ cells for SOD and $\text{nmol}/\text{min}/\text{mg}$ cells for GPx (SOD, superoxide dismutase; GPx, glutathione peroxidase).

decrease in absorbance was observed, suggesting the potential effect of α_2M on pesticide-treated cells, in contradiction with results obtained for group IV. Regarding antioxidant enzymes, SOD and GPx activities were quantified at 420 nm and 340 nm, respectively, and reported as units per milligram of cells for SOD and nanomoles per minute per milligram of cells for GPx³⁹. The results obtained for groups III, IV, and V suggest a cytoprotective role for α_2M against ROS-mediated toxicity induced by the pesticides. Because proteases and ROS work through a common mechanism of inflammation, like in diseases such as atherosclerosis⁴⁰, these findings also indicate the protective role of α_2M in preventing damage to various cellular species by ROS apart from trapping proteases.

CONCLUSIONS

Our data showed that the selected pesticides were potent neurotoxic agents, while α_2M decreased the cy-


totoxic effect of these pesticides. Hence, this study showed that α_2M modulates pesticide-induced ROS-based neurotoxicity based on different parameters. Thus, α_2M could possibly be used as a protective agent against neurotoxicity caused by ROS and pesticides, warranting further experimental studies.

ABBREVIATIONS

a2M: alpha-2-macroglobulin, **ALD:** aldicarb, **CPF:** chlorpyrifos, **DCFHDA:** dichlorodihydrofluorescein diacetate, **DLM:** deltamethrin, **GPx:** glutathione peroxidase, **H₂O₂:** hydrogen peroxide, **MDA:** malondialdehyde, melonaldehyde, **ROS:** reactive oxygen species, **SOD:** superoxide dismutase, **TBA:** thiobarbituric acid, **TBARS:** thiobarbituric acid reactive substances

ACKNOWLEDGMENTS

Research facilities provided by Department of Biochemistry, Aligarh Muslim University are gratefully



acknowledged.

AUTHOR'S CONTRIBUTIONS

SD: data collection, analysis and interpretation of results; drafted the manuscript.

HA: drafted the manuscript; critically revised and proofread the manuscript; approved the final version of manuscript. FHK: conception and design of work; approved the final version of manuscript. All authors read and approved the final manuscript.

FUNDING

SD is thankful to Indian Council of Medical Research (ICMR), New Delhi for an ICMR-SRF fellowship to support her research experiments.

AVAILABILITY OF DATA AND MATERIALS

Not applicable.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

CONSENT FOR PUBLICATION

Not applicable.

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

1. Abou-Donia MB, Lapidula DM. Mechanisms of organophosphorus ester-induced delayed neurotoxicity: type I and type II. Annual Review of Pharmacology and Toxicology. 1990;30(1):405–40. PMID: 2188574. Available from: [10.1146/annurev.pa.30.040190.002201](https://doi.org/10.1146/annurev.pa.30.040190.002201).
2. Imam A, Sulaiman NA, Oyewole AL, Chengetanai S, Williams V, Ajibola MI. Chlorpyrifos- and Dichlorvos-Induced Oxidative and Neurogenic Damage Elicits Neuro-Cognitive Deficits and Increases Anxiety-Like Behavior in Wild-Type Rats. Toxics. 2018;6(4):71. PMID: 30513797. Available from: [10.3390/toxics6040071](https://doi.org/10.3390/toxics6040071).
3. Dixit S, Zia MK, Siddiqui T, Ahsan H, Khan FH. Interaction of organophosphate pesticide chlorpyrifos with alpha-2-macroglobulin: biophysical and molecular docking approach. Journal of Immunoassay & Immunochemistry. 2021;42(2):138–53. PMID: 33086912. Available from: [10.1080/15321819.2020.1837161](https://doi.org/10.1080/15321819.2020.1837161).
4. Jokanović M. Neurotoxic effects of organophosphorus pesticides and possible association with neurodegenerative diseases in man: A review. Toxicology. 2018;410:125–31. PMID: 30266654. Available from: [10.1016/j.tox.2018.09.009](https://doi.org/10.1016/j.tox.2018.09.009).
5. Michaelis S, Burdon J, Howard CV. Aertoxic Syndrome: A New Occupational Disease? Public Health Panorama. 2017;3:198–211.
6. Lee JE, Park JH, Shin IC, Koh HC. Reactive oxygen species regulated mitochondria-mediated apoptosis in PC12 cells exposed to chlorpyrifos. Toxicology and Applied Pharmacology. 2012;263(2):148–62. PMID: 22714038. Available from: [10.1016/j.taap.2012.06.005](https://doi.org/10.1016/j.taap.2012.06.005).
7. Gupta SC, Mishra M, Sharma A, Balaji TGD, Kumar R, Mishra RK. Chlorpyrifos induces apoptosis and DNA damage in Drosophila through generation of reactive oxygen species. Ecotoxicology and Environmental Safety. 2010;73(6):1415–23. PMID: 20627310. Available from: [10.1016/j.ecoenv.2010.05.013](https://doi.org/10.1016/j.ecoenv.2010.05.013).

8. Chen R, Cui Y, Zhang X, Zhang Y, Chen M, Zhou T. Chlorpyrifos Induction of Testicular-Cell Apoptosis through Generation of Reactive Oxygen Species and Phosphorylation of AMPK. *Journal of Agricultural and Food Chemistry*. 2018;66(47):12455–70. PMID: 30378422. Available from: [10.1021/acs.jafc.8b03407](https://doi.org/10.1021/acs.jafc.8b03407).
9. Ki YW, Park JH, Lee JE, Shin IC, Koh HC. JNK and p38 MAPK regulate oxidative stress and the inflammatory response in chlorpyrifos-induced apoptosis. *Toxicology Letters*. 2013;218(3):235–45. PMID: 23416140. Available from: [10.1016/j.toxlet.2013.02.003](https://doi.org/10.1016/j.toxlet.2013.02.003).
10. Slotkin TA, Seidler FJ. Prenatal chlorpyrifos exposure elicits presynaptic serotonergic and dopaminergic hyperactivity at adolescence: critical periods for regional and sex-selective effects. *Reproductive Toxicology* (Elmsford, NY). 2007;23(3):421–7. PMID: 17267174. Available from: [10.1016/j.reprotox.2006.07.010](https://doi.org/10.1016/j.reprotox.2006.07.010).
11. Yang D, Howard A, Bruun D, Ajua-Alemanj M, Pickart C, Lein PJ. Chlorpyrifos and chlorpyrifos-oxon inhibit axonal growth by interfering with the morphogenic activity of acetylcholinesterase. *Toxicology and Applied Pharmacology*. 2008;228(1):32–41. PMID: 18076960. Available from: [10.1016/j.taap.2007.11.005](https://doi.org/10.1016/j.taap.2007.11.005).
12. Qiao D, Seidler FJ, Slotkin TA. Developmental neurotoxicity of chlorpyrifos modeled in vitro: comparative effects of metabolites and other cholinesterase inhibitors on DNA synthesis in PC12 and C6 cells. *Environmental Health Perspectives*. 2001;109(9):909–13. PMID: 11673119. Available from: [10.1289/ehp.01109909](https://doi.org/10.1289/ehp.01109909).
13. Eaton DL, Daroff RB, Atrupur H, Bridges J, Buffler P, Costa LG. Review of the toxicology of chlorpyrifos with an emphasis on human exposure and neurodevelopment. *Critical Reviews in Toxicology*. 2008;38(sup2):1–125. PMID: 18726789. Available from: [10.1080/10408440802272158](https://doi.org/10.1080/10408440802272158).
14. Yu F, Wang Z, Ju B, Wang Y, Wang J, Bai D. Apoptotic effect of organophosphorus insecticide chlorpyrifos on mouse retina in vivo via oxidative stress and protection of combination of vitamins C and E. *Experimental and Toxicologic Pathology*. 2008;59(6):415–23. PMID: 18222074. Available from: [10.1016/j.etp.2007.11.007](https://doi.org/10.1016/j.etp.2007.11.007).
15. Li Q, Kobayashi M, Kawada T. Chlorpyrifos induces apoptosis in human T cells. *Toxicology*. 2009;255(1-2):53–7. PMID: 18996166. Available from: [10.1016/j.tox.2008.10.003](https://doi.org/10.1016/j.tox.2008.10.003).
16. Park JH, Lee JE, Shin IC, Koh HC. Autophagy regulates chlorpyrifos-induced apoptosis in SH-SY5Y cells. *Toxicology and Applied Pharmacology*. 2013;268(1):55–67. PMID: 23352508. Available from: [10.1016/j.taap.2013.01.013](https://doi.org/10.1016/j.taap.2013.01.013).
17. Caughlan A, Newhouse K, Namgung U, Xia Z. Chlorpyrifos induces apoptosis in rat cortical neurons that is regulated by a balance between p38 and ERK/JNK MAP kinases. *Toxicological Sciences*. 2004;78(1):125–34. PMID: 14691213. Available from: [10.1093/toxsci/kfh038](https://doi.org/10.1093/toxsci/kfh038).
18. Yan D, Zhang Y, Liu L, Yan H. Pesticide exposure and risk of Alzheimer's disease: A systematic review and meta-analysis. *Sci Rep*. 2016;6:32222. PMID: 27581992. Available from: [10.1038/srep32222](https://doi.org/10.1038/srep32222).
19. Jiang X, Ganesan P, Rengarajan T, Choi DK, Arulselvan P. Cellular phenotypes as inflammatory mediators in Parkinson's disease: interventional targets and role of natural products. *Biomedicine and Pharmacotherapy*. 2018;106:1052–62. PMID: 30119171. Available from: [10.1016/j.biopha.2018.06.162](https://doi.org/10.1016/j.biopha.2018.06.162).
20. Sánchez-Santed F, Colomina MT, Hernández EH. Organophosphate pesticide exposure and neurodegeneration. *Cortex*. 2016;74:417–26. PMID: 26687930. Available from: [10.1016/j.cortex.2015.10.003](https://doi.org/10.1016/j.cortex.2015.10.003).
21. Dixit S, Zia MK, Siddiqui T, Ahsan H, Khan FH. Interaction of Human Alpha-2-Macroglobulin with Pesticide Aldicarb Using Spectroscopy and Molecular Docking. *Protein and Peptide Letters*. 2021;28(3):315–22. PMID: 32957873. Available from: [10.2174/0929866527666200921165834](https://doi.org/10.2174/0929866527666200921165834).
22. Blacker D, Wilcox MA, Laird NM, Rodes L, Horvath SM, Go RC. Alpha-2 macroglobulin is genetically associated with Alzheimer disease. *Nature Genetics*. 1998;19(4):357–60. PMID: 9697696. Available from: [10.1038/1243](https://doi.org/10.1038/1243).
23. Kovacs DM. α 2-macroglobulin in late-onset Alzheimer's disease. *Experimental Gerontology*. 2000;35(4):473–9. PMID: 10959035. Available from: [10.1016/S0531-5565\(00\)00113-3](https://doi.org/10.1016/S0531-5565(00)00113-3).
24. Varma VR, Varma S, An Y, Hohman TJ, Seddighi S, Casanova R. Alpha-2 macroglobulin in Alzheimer's disease: a marker of neuronal injury through the RCAN1 pathway. *Molecular Psychiatry*. 2017;22(1):13–23. PMID: 27872486. Available from: [10.1038/mp.2016.206](https://doi.org/10.1038/mp.2016.206).
25. Dixit S, Ahsan H, Khan FH. Pesticides and plasma proteins: unexplored dimensions in neurotoxicity. *International Journal of Pest Management*. 2021;p. 1–10. Available from: [10.1080/09670874.2021.1917725](https://doi.org/10.1080/09670874.2021.1917725).
26. Baek SH, Park T, Kang MG, Park D. Anti-Inflammatory Activity and ROS Regulation Effect of Sinapaldehyde in LPS-Stimulated RAW 264.7 Macrophages. *Molecules* (Basel, Switzerland). 2020;25(18):4089. PMID: 32906766. Available from: [10.3390/molecules25184089](https://doi.org/10.3390/molecules25184089).
27. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*. 1983;65(1-2):55–63. PMID: 6606682. Available from: [10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4).
28. LeBel CP, Ischiropoulos H, Bondy SC. Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chemical Research in Toxicology*. 1992;5(2):227–31. PMID: 1322737. Available from: [10.1021/tx00026a012](https://doi.org/10.1021/tx00026a012).
29. Wang H, Joseph JA. Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radical Biology & Medicine*. 1999;27(5-6):612–6. PMID: 10490282. Available from: [10.1016/S0891-5849\(99\)00107-0](https://doi.org/10.1016/S0891-5849(99)00107-0).
30. Gay C, Gebicki JM. A critical evaluation of the effect of sorbitol on the ferric-xylenol orange hydroperoxide assay. *Analytical Biochemistry*. 2000;284(2):217–20. PMID: 10964403. Available from: [10.1006/abio.2000.4696](https://doi.org/10.1006/abio.2000.4696).
31. Hasanzadeh D, Mahdavi M, Dehghan G, Charoudeh HN. Farnesiferol C induces cell cycle arrest and apoptosis mediated by oxidative stress in MCF-7 cell line. *Toxicology Reports*. 2017;4:420–6. PMID: 28959668. Available from: [10.1016/j.toxrep.2017.07.010](https://doi.org/10.1016/j.toxrep.2017.07.010).
32. Marklund S, Marklund G. Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*. 1974;47(3):469–74. PMID: 4215654. Available from: [10.1111/j.1432-1033.1974.tb03714.x](https://doi.org/10.1111/j.1432-1033.1974.tb03714.x).
33. Flohé L, Günzler WA. Assays of glutathione peroxidase. *Methods in Enzymology*. 1984;105:114–21. PMID: 6727659. Available from: [10.1016/S0076-6879\(84\)05015-1](https://doi.org/10.1016/S0076-6879(84)05015-1).
34. Aktar MW, Sengupta D, Chowdhury A. Impact of pesticides use in agriculture: their benefits and hazards. *Interdisciplinary Toxicology*. 2009;2(1):1–12. PMID: 21217838. Available from: [10.2478/v10102-009-0001-7](https://doi.org/10.2478/v10102-009-0001-7).
35. Andersson H, Tago D, Treich N. A review of evidence on health effects valuation of risks and benefit-cost analysis. *Toulouse School of economics. Pestic Health*. 2014;24:203–95.
36. de Oliveira MR, Peres A, Ferreira GC, Schuck PF, Bosco SM. Carnosic Acid Affords Mitochondrial Protection in Chlorpyrifos-Treated Sh-Sy5y Cells. *Neurotoxicity Research*. 2016;30(3):367–79. PMID: 27083155. Available from: [10.1007/s12640-016-9620-x](https://doi.org/10.1007/s12640-016-9620-x).
37. Green DR, Reed JC. Mitochondria and apoptosis. *Science*. 1998;281(5381):1309–12. PMID: 9721092. Available from: [10.1126/science.281.5381.1309](https://doi.org/10.1126/science.281.5381.1309).
38. Keller A, Mohamed A, Dröse S, Brandt U, Fleming I, Brandes RP. Analysis of dichlorodihydrofluorescein and dihydrocalcein as probes for the detection of intracellular reactive oxygen species. *Free Radical Research*. 2004;38(12):1257–67. PMID: 15763950. Available from: [10.1080/10715760400022145](https://doi.org/10.1080/10715760400022145).
39. Wasi S, Tabrez S, Ahmad M. Toxicological effects of major environmental pollutants: an overview. *Environmental Monitor-*

- ing and Assessment. 2013;185(3):2585–93. PMID: [22763655](#). Available from: [10.1007/s10661-012-2732-8](#).
40. Slack MA, Gordon SM. Protease Activity in Vascular Disease. Arteriosclerosis, Thrombosis, and Vascular Biology. 2019;39(10):e210–8. PMID: [31553665](#). Available from: [10.1161/ATVBAHA.119.312413](#).